

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/113007/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Beltrami, Cristina, Simpson, Kate, Jesky, Mark, Wonnacott, Alexa ORCID: <https://orcid.org/0000-0002-0968-4248>, Carrington, Christopher, Holmans, Peter ORCID: <https://orcid.org/0000-0003-0870-9412>, Newbury, Lucy, Jenkins, Robert ORCID: <https://orcid.org/0000-0001-8500-9044>, Ashdown, Thomas, Dayan, Colin ORCID: <https://orcid.org/0000-0002-6557-3462>, Satchell, Simon, Corish, Peter, Cockwell, Paul, Fraser, Donald ORCID: <https://orcid.org/0000-0003-0102-9342> and Bowen, Timothy ORCID: <https://orcid.org/0000-0001-6050-0435> 2018. Association of elevated urinary miR-126, miR-155, and miR-29b with diabetic kidney disease. American Journal of Pathology 188 (9) , pp. 1982-1992. 10.1016/j.ajpath.2018.06.006 file

Publishers page: <http://dx.doi.org/10.1016/j.ajpath.2018.06.006>
<<http://dx.doi.org/10.1016/j.ajpath.2018.06.006>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



**Association of Elevated Urinary
miR-126, miR-155 and miR-29b with Diabetic Kidney Disease**

Cristina Beltrami,^{*} Kate Simpson,^{*†} Mark Jesky,[‡] Alexa Wonnacott,^{*}
Christopher Carrington,^{*} Peter Holmans,[§] Lucy Newbury,^{*†} Robert Jenkins,^{*}
Thomas Ashdown,^{*} Colin Dayan,[¶] Simon Satchell,^{||} Peter Corish,^{**}
Paul Cockwell,[‡] Donald Fraser,^{*†‡‡} and Timothy Bowen^{*†‡‡§§}

^{*} Wales Kidney Research Unit, Division of Infection and Immunity, School of Medicine, College of Biomedical and Life Sciences, Cardiff University, Heath Park Campus, Cardiff CF14 4XN, UK

[†] Cardiff Institute of Tissue Engineering and Repair, Cardiff University, Museum Place, Cardiff CF10 3BG, UK

[‡] Department of Renal Medicine, Queen Elizabeth Hospital Birmingham, Mindelsohn Way, Birmingham B15 2GW, UK

[§] MRC Centre for Neuropsychiatric Genetics and Genomics, Division of Psychological Medicine and Clinical Neurosciences, School of Medicine, College of Biomedical and Life Sciences, Cardiff University, Maindy Road, Cathays, Cardiff CF24 4HQ, UK

[¶] Diabetes Research Group, Division of Infection and Immunity, School of Medicine, College of Biomedical and Life Sciences, Cardiff University, Heath Park Campus, Cardiff CF14 4XN, UK

^{||} Bristol Renal, Bristol Medical School, University of Bristol, Dorothy Hodgkin Building, Whitson Street, Bristol BS1 3NY, UK

^{**} BBI Group, The Courtyard, Ty Glas Avenue, Cardiff CF14 5DX, UK

^{‡‡} These authors contributed equally to this work

Running title: Urinary miRNAs in Diabetic Nephropathy

^{§§} To whom correspondence should be addressed:

Timothy Bowen, Wales Kidney Research Unit, School of Medicine, College of Biomedical and Life Sciences, Cardiff University, Heath Park Campus, Cardiff CF14 4XN, UK. Tel.: 44-29-2074-8389; Fax: 44-29-2074-8470; E-mail: bowent@cf.ac.uk

Counts Abstract: 215 words; Text: 15 pages; Tables: 3; Figures: 4

Support This is a publication of independent research funded by the National Institute for Health Research (NIHR) Invention for Innovation (i4i) Programme (Grant Reference no. II-LA-0712-20003). The Principal Investigator for the grant is TB. The views expressed in this paper are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health, UK. The authors also acknowledge support from Kidney Research UK Project Grant Awards RP44/2014 (TB) and IN4/2013 (SS). The JABBS Foundation funded collection of the Renal Impairment in Secondary Care (RIISC) cohort (PC). The Wales Kidney Research Unit is funded by core support from Health and Care Research Wales (DJF).

Declaration of interests CD: Novo Nordisk Advisor, Sanofi Genzyme and AstraZeneca services (2015-2017); SS: Boehringer Ingelheim Travel Support and UCB UK Grant Support (2013-2015); PCor: Life Sciences Bridging Fund Wales Consultancy (2015-2017); PC: JABBS Foundation Grant Support (2014-2015); DF and TB: Patent application PCT/GB2017/050195: Kidney disease diagnostic (2017).

1 **ABSTRACT**

2 Effective diabetic kidney disease (DKD) biomarkers remain elusive, and urinary
3 microRNAs (miRNAs) represent a potential source of novel non-invasive disease
4 sentinels. We profiled 754 miRNAs in pooled urine samples from DKD patients (n = 20),
5 detecting significantly increased miR-126, miR-155 and miR-29b compared to controls
6 (n = 20). These results were confirmed in an independent cohort of 89 DKD patients,
7 62 diabetic patients without DKD and 41 controls: miR-126 (2.8-fold increase;
8 p<0.0001), miR-155 (1.8-fold; p<0.001) and miR-29b (4.6-fold; p = 0.024). Combined
9 receiver operating characteristic curve analysis resulted in an area under the curve of
10 0.8. A relative quantification threshold equivalent to 80% sensitivity for each miRNA
11 gave a positive signal for 48% of DKD patients compared to 3.6% of diabetic patients
12 without DKD. Laser capture microdissection of renal biopsies followed by RT-qPCR
13 detected miR-155 in glomeruli, proximal and distal tubules, while miR-126 and miR-
14 29b were most abundant in glomerular extracts. Subsequent experiments showed
15 miR-126 and miR-29b enrichment in glomerular endothelial cells (GEnCs) compared to
16 podocytes, proximal tubular epithelial cells and fibroblasts. Significantly increased miR-
17 126 and miR-29b were detected in GEnC conditioned medium in response to tumour
18 necrosis factor-alpha and transforming growth factor-beta 1, respectively. Our data
19 reveal an altered urinary miRNA profile associated with DKD and link these variations
20 to miRNA release from GEnCs.

21

22 **Keywords:** microRNAs, urine, biomarker, diabetic kidney disease, chronic kidney
23 disease

24

25 **Abbreviations:**

26 ACR albumin:creatinine ratio

27 CKD chronic kidney disease

28 DKD diabetic kidney disease

29 eGFR estimated glomerular filtration rate

30 GEnC glomerular endothelial cell

31 KDIGO kidney disease: improving global outcomes

32 NKF KDOQI National Kidney Foundation kidney disease outcomes quality initiative

33 LCM laser capture microdissection

34 MDRD modification of diet in renal disease

35 miRNA microRNA

36 PTC proximal tubular epithelial cell

37 RIISC renal impairment in secondary care study

38

39

40 **Introduction**

41 Recent estimates suggest that 1 in 12 of the global population suffers from diabetes
42 mellitus, approximately 40 % of those affected will go on to develop diabetic kidney
43 disease (DKD).¹ DKD is the leading cause of end-stage renal disease and predisposing
44 factors include genetic causes, ethnicity, hyperglycaemia, insulin resistance,
45 intraglomerular hypertension and hyperfiltration.^{2,3}

46 Hyperglycaemia results in numerous deleterious consequences including
47 upregulated cytokine synthesis, renin-angiotensin system activation, generation of
48 advanced glycation end products and reactive oxygen species, and increased protein
49 kinase C activity.^{4,5} Nitric oxide and NF-κB pathway-driven loss of endothelial and
50 vascular modulation have been implicated in insulin resistance, and early DKD may
51 be associated with insulin signaling defects specific to the podocyte.⁶ These insults
52 result in loss of glomerular filtration rate and ultimately to renal failure from
53 mesangial hyperexpansion, nodular glomerulosclerosis and tubulointerstitial
54 fibrosis.⁷

55 Detection of urinary microalbuminuria currently forms the basis of DKD
56 progression monitoring, varying from normal mean albuminuria values around 10
57 mg/day to a diagnosis of microalbuminuria at 30–300 mg/day and macroalbuminuria
58 above 300 mg/day.⁸ Prognosis is complicated, since not all microalbuminuric
59 patients progress to overt nephropathy. A number of novel biomarkers have been
60 assessed for utility in DKD but none are being used as routine clinical markers, and
61 they may lack specificity and sensitivity to predict individual DKD patient outcomes.
62 In light of the above, novel markers that can discriminate aetiology, progression
63 and/or response to treatment remain highly desirable.

MicroRNAs (miRNAs) are ubiquitously-expressed short noncoding RNAs that regulate the expression of most protein coding genes in the human genome, and detection of miR-192, miR-194, miR-215, miR-216, miR-146a, miR-204 and miR-886 is elevated in the kidney.⁹ Urinary miRNAs represent a highly promising novel source of non-invasive biomarkers that are stabilised via argonaute 2 protein/exosome association and are rapidly and precisely detected by RT-qPCR.¹⁰

Reports have suggested a role for miRNAs in the pathology of DKD,^{11,12} including previous work from this laboratory showing decreased miR-192 in biopsies from late-stage DKD patients with diminished renal function.¹³ However, comparatively little is known about the abundance of urinary miRNAs in DKD patients.

We hypothesised that alterations in urinary miRNA profiles would be associated with DKD. We identified candidate DKD biomarkers by comparing miRNA profiles in urine samples from a patient discovery cohort with those from unaffected controls. Selected candidates were then measured in a larger, independent cohort. Subsequently, laser capture microdissection of renal biopsies and *in vitro* cell culture were used to investigate the sources of our candidate urinary miRNA DKD biomarkers with respect to nephron domain and cell type.

Materials and Methods

Study Participants

DKD was defined in accordance with the National Kidney Foundation Kidney Disease Outcomes Quality Initiative (NKF KDOQI) Clinical Practice Guidelines and Clinical Practice Recommendations for Diabetes and Chronic Kidney Disease (CKD).¹⁴ Accordingly, CKD should be attributable to diabetes in the presence of macroalbuminuria (in the absence of urinary infection), or in the presence of microalbuminuria with concomitant diabetic retinopathy, or in type 1 diabetes of at least 10 years duration.¹⁴ The initial profiling study cohort of 20 DKD patients and 20 healthy controls was obtained from the Wales Kidney Research Tissue Bank, University Hospital of Wales, Cardiff. The DKD group was predominantly male (85%), mean age 72 years (SD +/- 8.7). DKD patients were CKD stage 3-5 (pre-dialysis), with mean eGFR of 29 ml/min/1.73m² (SD +/- 8.5) and a mean urinary Albumin:Creatinine ratio (ACR) of 13.5 mg/mmol (SD +/- 14.5). The control group (n = 20) in the profiling cohort were 50% male, mean age 47 years (SD +/- 11.0) with no microalbuminuria (ACR<3 mg/mmol). For further details on ACR categories see Table 1.

The confirmation cohort was drawn from two secondary care facilities: the Wales Kidney Research Tissue Bank (as above) and the Renal Impairment In Secondary Care (RIISC) study, University Hospital of Birmingham, UK.¹⁵ 89 patients with DKD, including 3 patients with type 1 diabetes, and 41 healthy controls were recruited across the two sites. An additional control group of 62 diabetics without DKD were recruited from Cardiff, including 17 patients with type 1 diabetes. Ethical approval

was granted by the Wales Kidney Research Tissue Bank Governance Committee and the South Birmingham Local Research Ethics Committee, respectively.

Patient demographics and clinical parameters are shown in Table 1. All patients were recruited from specialist nephrology and diabetes care services at the two sites during the period spanning autumn 2010 to autumn 2013. DKD patients from the RIISC study cohort were predominantly advanced nephropaths as per RIISC protocol inclusion criteria: briefly, patients with CKD stages 4-5 (pre-dialysis), or CKD stage 3 and accelerated progression and/or proteinuria as defined by the UK National Institute for Health and Care Excellence 2008 CKD guideline for secondary care review. The diabetic patient control group all had a diagnosis of diabetes by standard American Diabetes Association criteria,¹⁶ but without evidence of DKD (i.e. not fulfilling the KDOQI criteria).

At initial clinic visit, renal function was recorded using estimated glomerular filtration rate (eGFR), calculated using the modification of diet in renal disease (MDRD) equation¹⁷. Urine samples were aliquoted for albumin:creatinine ratio (ACR) assessment and for RNA extraction (see below). ACR cut-offs for disease severity were defined as per Kidney Disease: Improving Global Outcomes (KDIGO) 2012 guidelines.¹⁸

Urine Collection, RNA Isolation and RT-qPCR analysis

Urine samples were collected and RNA extraction from 350 µl of urine, generation of cDNA from equal volumes of RNA extracts and RT-qPCR were then carried out as

described in detail elsewhere.¹⁰ TaqMan assays (Thermo Fisher Scientific, Paisley, UK) used in this study were: hsa-miR-29b-3p (ID 000413); hsa-miR-126-3p (ID 002228); hsa-miR-155-5p (ID 002623); hsa-miR-191-5p (ID 002299). Relative quantities were calculated using the $2^{-\Delta\Delta Ct}$ method, and miRNA expression was normalized to hsa-miR-191-5p.¹⁰

MiRNA profiling by TaqManArray Human MicroRNA Cards

Urinary miRNAs were reverse transcribed using the Megaplex Primer Pools (Human Pools A v2.1 and B v3.0, Thermo Fisher Scientific) with a predefined pool of 381 reverse transcription (RT) primers for each Megaplex Primer Pool. A fixed volume of 3 µl of RNA solution was used as input in each RT reaction, and RT reactions were performed according to the manufacturer's recommendations. RT reaction products were amplified using Megaplex PreAmp Primers (Primers A v2.1 and B v3.0, Thermo Fisher Scientific), the samples were then diluted to a final volume of 100 µl and control subject and DKD patient products were pooled as follows.

To exclude the possibility that gender, age and eGFR status had extreme effects on miRNA expression profiles, the following pooling strategy was followed. Control pool (CP)1: urine samples from 5 females of average age 44.8 years; CP2: 5 females, 57.6; CP3: 5 males, 35.2; CP4, 5 males, 53.2. Patient Pool (PP)1: urine samples from 5 CKD3 patients with an eGFR between 43.3 and 36 mL/min per 1.73m²; PP2: 5 stage 3 patients, 35 - 31; PP3: 5 stage 4/5 patients, 27.3 - 23; PP4: 5 stage 4/5 patients, 22 - 12.9.

TaqManArray Human MicroRNA Cards A v2.1 and B v3.0 (Thermo Fisher Scientific) were used to quantify 754 human miRNAs. Each array included 377 test miRNAs, 3 endogenous controls and a negative control. Quantitative (q)PCR was carried out on an Applied Biosystems 7900HT thermo cycler (Thermo Fisher Scientific) using the manufacturer's recommended program.

Laser Capture Microdissection (LCM) from Renal Biopsy Samples

Glomeruli, proximal tubular and distal tubular profiles were microdissected from 6- μ m sections obtained from five FFPE archived renal biopsies from unaffected people using the Arcturus Pixcell Ite infrared laser enabled LCM system (Thermo Fisher Scientific).

Cell Culture

Human conditionally immortalised glomerular endothelial cell (GEnC) and human podocyte (ATC) cell lines were propagated at 33°C as described previously.^{19,20} After 5 (GEnC) and 14 (ATC) days, cells were transferred to 37°C incubation to inactivate the SV40 T antigen and permit differentiation, prior to experimental use. Where stated, GEnCs were growth arrested for 24 h and then treated with TNF- α (10 ng/mL) or TGF- β 1 (1 ng/mL) at either normoglycaemic (5 mM) or hyperglycaemic (25 mM) D-glucose concentrations for 24 h. Proximal tubular epithelial cell (PTC) line HK-2²¹ and fibroblast²² cultures were maintained as described elsewhere. Cells and culture medium obtained from each well were used for RNA extraction as described above.

178 **Statistical analysis**

179 MiRNA profiling data were analysed using Thermo Fisher Scientific's DataAssist
180 Software (version 3.01), NormFinder Software ([http://moma.dk/normfinder-](http://moma.dk/normfinder-software)
181 software; last access 21/02/18) and GraphPad Prism 6 (version 6.0d). Pearson
182 Correlation Coefficients was used to detect clusters of similarity in miRNA threshold
183 cycle values between each pool group in patients, and between each pool group in
184 controls. To identify a suitable reference gene for the normalization of miRNA
185 expression in this study, the NormFinder algorithm was applied to the expression
186 data obtained from the Human TaqMan miRNA Arrays. Analysis comparing miRNA
187 levels between subjects with DKD and controls was carried out using GraphPad Prism
188 version 6 version 6.0d. Values for p below 0.05 were considered statistically
189 significant. MiRNA profiling data sets can be found in Gene Expression Omnibus
190 (<https://www.ncbi.nlm.nih.gov/geo>; accession number GSE114477).

191

Results

Altered urinary miRNA detection in DKD patients

To select candidate miRNAs that may act as DKD biomarkers, we first compared data from unbiased expression profiling of 754 miRNAs in urine samples from 20 DKD patients and 20 unaffected controls. Analyses were performed on 4 patient and 4 control pools, each composed of urine samples from 5 individuals as recommended by Zhang and colleagues.²³ Samples were pooled prior to profiling to minimize the contribution of subject to subject variation and to make substantive features easier to find, and thereby identify biomarkers common across individuals.²⁴ Previous analysis suggested that 40 individuals might optimally be pooled across 8 arrays,²³ which was our chosen pooling approach.

In Figure 1A the 12 data points in the upper right quadrant of the plot represent those miRNAs for which statistically significant fold-change increases were detected in patient urine compared to control samples, the 35 points in the upper left quadrant the corresponding downregulated miRNAs. The fold-change data for these 47 miRNAs are summarised in Figure 1B, and the 8 miRNAs exhibiting >5-fold change were subsequently selected as potential candidate biomarkers for further analysis.

Specific RT-qPCR assays were then used to analyse these miRNAs in each component urine sample pooled for profiling analysis. Statistically significant differences in miRNA detection between DKD patient and control urine samples were replicated for miR-126 (4.3-fold increase; $p = 0.0087$), miR-155 (22.9-fold; $p = 0.0024$) and miR-29b (4.9-fold; $p = 0.0002$) (Figure 1, C-E).

Elevated urinary miR-126, miR-155 and miR-29b detection in an independent DKD patient cohort

To test the above findings, miR-126, miR-155 and miR-29b were quantified in samples from an independent cohort of patients with established DKD from the Renal Impairment in Secondary Care Study (RIISC).¹⁵ Samples from 89 patients meeting the criteria established in the UK National Institute for Health and Care Excellence 2008 criteria were available. An additional cohort of 62 patients with diabetes mellitus but without proteinuria or other evidence of DKD were included, as were samples from 41 people without evidence of diabetes or DKD (Table 1). We included diabetes patients without DKD as a third group in this analysis to identify DKD-specific miRNA detection changes and not purely hyperglycaemia-driven effects from our profiling comparison of DKD patients with control individuals.

Significant differences were again seen between DKD patients and controls for miR-126 (2.8-fold increase; $p < 0.0001$; Figure 2A), miR-155 (1.8-fold; $p < 0.001$; Figure 2B) and miR-29b (4.6-fold; $p = 0.024$; Figure 2C). Comparison of DKD patients with diabetic patients without DKD was statistically significant for miR-126 (3.1-fold increase; $p < 0.0001$) and miR-155 (1.6-fold; $p = 0.024$) with a trend to increased miR-29b (4.1-fold; $p = 0.121$) (Figure 2A-C).

RT-qPCR data for all 3 miRNAs were used to compare DKD patients and diabetic patients without DKD in the combined receiver operating characteristic (ROC) curve analysis shown in Figure 2D, giving an area under the curve (AUC) of 0.80. To analyse the contributions of each miRNA to the above ROC curve, individual specificity and likelihood ratios were calculated for relative expression (RQ) values

equivalent to a sensitivity of 80%.^{25,26} Data displayed in Table 2 illustrate the magnitude of corresponding specificity values was miR-126 > miR-155 > miR-29b, and that combined miRNA data resulted in a $\geq 6.5\%$ increase in specificity and likelihood ratio compared with individual miRNAs. These RQ data were then used as consecutive threshold values to discriminate between DKD and diabetic patients without DKD (D in Table 3) from the independent cohort. The discriminatory order was miR-29b (DKD/D = 5.62) > miR-126 (3.48) > miR-155 (2.23), and RQ values exceeding all 3 thresholds were obtained for 48.0% of DKD patients compared with 3.6% of diabetic patients without DKD (Table 3).

Laser capture microdissection shows increased glomerular abundance of miR-126 and miR-29b that is replicated in GEnC culture

Previous reports have linked changes in miRNA expression to DKD pathology, but have focused on whole tissue studies. For example, we showed association of decreased miR-192 expression with disease progression in DKD biopsies by *in situ* hybridisation.¹³

In the present study we used laser capture microdissection (LCM) to isolate glomeruli, proximal and distal tubules (Figure 3A) from histologically normal formalin-fixed, paraffin-embedded (FFPE) renal biopsy samples, and analysed miR-126, miR-155 and miR-29b expression by RT-qPCR. In Figure 3B, a typical CD10-stained FFPE biopsy section is seen before and after LCM to isolate glomeruli, proximal and distal renal tubules. MiR-126, miR-155 and miR-29b were detected in extracts from all three nephron regions (Figure 3, C-E). Increased glomerular

abundances were observed for miR-126 (Figure 3C) and miR-29b (Figure 3E), while miR-155 was most abundant in the distal tubule (Figure 3D).

Conclusions regarding nephron region-specific miRNA expression from the above analyses are inherently limited, however, since tissue extracts are subject to trace contamination by cells from other nephron domains. Therefore, cellular miRNA localisation within each nephron region was subsequently investigated by RT-qPCR analysis of podocyte and endothelial cell (GEnC) cultures from the glomerulus, renal proximal tubular epithelial cells (PTC) and fibroblasts. Detection of miR-126 was significantly higher in GEnCs compared with other cell types (Figure 3F). Most miR-155 was detected in PTCs and least in GEnCs (Figure 3G), while miR-29b was most abundant in GEnCs (Figure 3H).

GEnC release of miR-126 and miR-29b in an *in vitro* model of hyperglycaemia is driven by TNF- α and TGF- β 1, respectively

The above data localized the majority of miR-126 and miR-29b expression to the GEnC. We next sought stimuli by which miRNAs are released into the glomerular ultrafiltrate, and hence the urine. Data from animal models of diabetes show increased glomerular and PTC TNF- α expression, and renoprotective effects of TGF- β inhibitors have also been reported.^{27,28} GEnC expression of our candidate miRNAs was thus analysed *in vitro* in response to TNF- α and TGF- β 1 in normoglycaemia and hyperglycaemia (Figure 4, A-F).

The presence of TNF- α led to significantly increased miR-126 detection in GEnC conditioned medium at 5 mM and 25 mM D-glucose (Figure 4B), a pattern also seen for miR-29b following TGF- β 1 addition (Figure 4D). These cytokines did not

increase GEnC expression of miR-126 (Figure 4A), or miR-29b (Figure 4C), a pattern consistent with increased release, but not expression, of miRNAs.

No significant changes in miR-155 were detected in response to elevated D-glucose with either cytokine, and data for TNF- α are shown (Figure 4, E and F). Similarly, changes in miR-126 following TGF- β 1 addition, and for miR-29b in the presence of TNF- α , were not observed (data not shown). Elevated D-glucose alone did not change miRNA expression in GEnCs or conditioned medium (Figure 4, A-F).

Discussion

Diabetic kidney disease (DKD) is the leading cause of kidney failure requiring renal replacement therapy worldwide, but effective methods to identify and halt progression of disease-specific pathophysiological changes remain elusive. Current effective interventions such as control of blood glucose and blood pressure are challenging to achieve, costly and time intensive. Existing tests track DKD from diabetic diagnosis to kidney failure, but do not allow accurate prognosis for the individual patient. In addition, the absence of treatment response biomarkers hinders development of emerging DKD therapies. There is thus an unmet need for additional DKD biomarkers to target intervention and follow response to therapy.

In this study we set out to identify urinary miRNA DKD biomarkers. Increased detection of miR-126, miR-155 and miR-29b was observed in the urine of DKD patients in comparison with both unaffected individuals and diabetic patients without DKD. MiRNA localization and release studies further suggested specific release of miR-126 and miR-29b from GEnCs. This raised the possibility that urinary miRNA quantification might provide data on ongoing pathological processes, and so aid patient stratification and measurement of response to therapy.

Urinary miRNA biomarkers have several potential significant advantages over circulating miRNAs for adoption into existing treatment pathways alongside current biomarkers, including speed and cost of non-invasive sample access.²⁹ However, few urinary miRNA DKD biomarker data have so far been reported. Previous studies have focused on circulating miRNAs, and have generated conflicting data with respect to association of miR-126 with diabetes mellitus and/or DKD. A recent cross-sectional analysis of type 2 diabetes mellitus patients found a negative association with

plasma miR-126,³⁰ and similar findings have been reported for type 1 diabetes mellitus and all complications.³¹ By contrast, miR-126 detection did not change in whole blood from type 2 diabetes mellitus patients and control subjects, but decreased in DKD patient samples.³² Furthermore, no change in plasma miR-126 was observed in a study of paediatric type 1 diabetic patients.³³ These analyses provide inconsistent data for the biomarker utility of circulating miR-126, in contrast to the significant and reproducible increases we detected in miR-126, miR-155 and miR-29b in DKD patient urine in the present study.

The DKD-specific alterations in urinary miRNA profiles detected in this study may have functional significance. Our *in vitro* analyses localized miR-126 and miR-29b principally to the GEnC, with miR-155 expression distributed evenly across the nephron. Glomerular endothelial localization of miR-126 may reflect the role of this transcript in vascular regulation. Targeted mouse miR-126 deletion resulted in vascular abnormalities by removing inhibition of sprouty-related EVH1 domain-containing protein 1 expression, thereby enhancing vascular endothelial growth factor (VEGF) function.³⁴ A role in DKD pathology for VEGFA signalling between GEnCs and podocytes has been proposed.³⁵ In addition, miR-126 repression of vascular cell adhesion molecule 1 expression in human umbilical vein endothelial cells regulates their response to pro-inflammatory adhesion molecules.³⁶ MiR-126 has also been implicated in the heterogenic inflammatory response of renal microvascular endothelial cells.³⁷

Increased expression of miR-155 has been observed in DKD patient renal biopsies, in close correlation with increased serum creatinine.³⁸ Furthermore, miR-155 deficiency attenuated renal damage and IL-17 expression was downregulated in

streptozotocin-induced DKD mice.³⁹ Together with miR-126, miR-155 has been implicated in multiple forms of vascular remodelling and associated with cardiovascular disease.⁴⁰

Decreased miR-29b has been reported in early and advanced animal models of diabetic renal fibrosis.⁴¹ Chen and colleagues found that loss of renal miR-29b in db/db mice led to increased albuminuria, TGF- β -mediated fibrosis and immune injury, while restored miR-29b expression inhibited renal injury.⁴² Indeed, while we have focused on upregulated miRNAs in this study, we acknowledge the potential importance of miRNA downregulations that we detected.

In the present study we localized miR-29b to the glomerular endothelium. Reduction of collagen and laminin synthesis has been reported following forced miR-29b expression in human corneal endothelial cells.⁴³ In apoE knockout mice, miR-29b induced aortic endothelial permeability in response to a high fat diet, and brought about aortic apoptosis by direct targeting of melatonin receptor mt1.⁴⁴ In addition, upregulated miR-29b expression has been observed in human umbilical vein endothelial cells exposed to hyperglycaemia.⁴⁵

The cytokine-driven release from GEnCs observed for miR-126 (TNF- α) and miR-29b (TGF- β 1) reported here suggests that these cells may be the principal source of elevated urinary miR-126 and miR-29b detected in DKD. We speculate that this constitutes evidence for disease-related signalling down the nephron that will be interesting to test in future studies. Indeed, we have demonstrated association of urinary miRNAs with exosomes¹⁰ and exosomal transport, which might facilitate passage of miRNAs through the nephron, has been reported for all three candidate biomarker miRNAs.

365 Exosome-mediated release of miR-126 from CD34⁺ peripheral blood
366 mononuclear cells is proangiogenic, and decreased miR-126 was detected in
367 elevated glucose cell culture and diabetic patients.⁴⁶ MiR-155 is depleted in urinary
368 exosomes from microalbuminuric type 1 diabetes mellitus patients.⁴⁷ Endogenous
369 miR-29b, spontaneously released from beta-cells within exosomes, stimulates TNF-
370 α secretion from spleen cells isolated from diabetes-prone NOD mice *in vitro*.⁴⁸

371 In summary, we have used unbiased profiling approaches to identify a urinary
372 miRNA signature associated with DKD, and have subsequently confirmed increased
373 miR-126, miR-155 and miR-29b in an independent patient cohort. MiR-126 and miR-
374 29b were identified as enriched in GEnCs, and released from these cells in response
375 to DKD-related cytokines. Urinary miR-126, miR-155 and miR-29b are therefore
376 promising DKD biomarkers, and the potential pathological significance of miR-126
377 and miR-29b release from GEnCs merits further evaluation.

378

Acknowledgments

We thank control subjects and patients for the donation of urine samples including those samples kindly provided by co-authors Dr Mark D. Jesky and Professor Paul Cockwell (Queen Elizabeth Hospital Birmingham, Birmingham, UK).

C.B. and K.S. performed the experimental work, generated and analyzed data, and helped draft the manuscript. A.W., C.C., L.N., R.J. and T.A. performed the experimental work, generated and analyzed data. M.J., P.H., C.D., S.S., P.Cor and P.C. discussed elements of experimental design and/or cohort composition. D.F. and T.B. designed the research. T.B. wrote the manuscript, which was edited by D.F. then amended and approved by each author.

References

1. Jha V, Garcia-Garcia G, Iseki K, Li Z, Naicker S, Plattner B, Saran R, Wang AY, Yang CW: Chronic kidney disease: global dimension and perspectives. *Lancet* 2013, 382:260–272
2. Cowie CC, Port FK, Wolfe RA, Savage PJ, Moll PP, Hawthorne VM: Disparities in incidence of diabetic end-stage renal disease according to race and type of diabetes. *N Engl J Med* 1989, 321:1074–1079
3. Regele F, Jelencsics K, Shiffman D, Pare G, McQueen MJ, Mann JF, Oberbauer R: Genome-wide studies to identify risk factors for kidney disease with a focus on patients with diabetes. *Nephrol Dial Transplant* 2015, 30:26–34
4. Makita Z, Radoff S, Rayfield EJ, Yang Z, Skolnik E, Delaney V, Friedman EA, Cerami A, Viassara H: Advanced glycosylation end products in patients with diabetic nephropathy. *N Engl J Med* 1991, 325:836–842
5. Schrijvers BF, De Vriese AS, Flyvbjerg A: From hyperglycemia to diabetic kidney disease: the role of metabolic, hemodynamic, intracellular factors and growth factors/cytokines. *Endocr Rev* 2004, 25:971–1010
6. Coward RJ, Saleem MA: Podocytes as a target of insulin. *Curr Diabetes Rev* 2011, 7:22–27
7. Rodriguez-Iturbe B, Johnson RJ, Herrera-Acosta J: Tubulointerstitial damage and progression of renal failure. *Kidney Int Suppl* 2005, 99:S82–S86
8. Parving H-H, Persson F, Rossing P: Microalbuminuria: A parameter that has changed diabetes care. *Diab Res Clin Pract* 2015, 107:1-8
9. Landgraf P, Rusu M, Sheridan R, Sewer A, Iovino N, Aravin A, Pfeffer S, Rice A, Kamphorst AO, Landthaler M, Lin C, Socci ND, Hermida L, Fulci V, Chiaretti S,

414 Foà R, Schliwka J, Fuchs U, Novosel A, Müller RU, Schermer B, Bissels U,
415 Inman J, Phan Q, Chien M, Weir DB, Choksi R, De Vita G, Frezzetti D,
416 Trompeter HI, Hornung V, Teng G, Hartmann G, Palkovits M, Di Lauro R,
417 Wernet P, Macino G, Rogler CE, Nagle JW, Ju J, Papavasiliou FN, Benzing T,
418 Lichter P, Tam W, Brownstein MJ, Bosio A, Borkhardt A, Russo JJ, Sander C,
419 Zavolan M, Tuschl T: A mammalian microRNA expression atlas based on small
420 microRNA sequencing. *Cell* 2007, 129:1404-1414

421 10. Beltrami C, Clayton A, Newbury LJ, Corish C, Jenkins RH, Phillips AO, Fraser DJ,
422 Bowen T: Stabilization of urinary microRNAs by association with exosomes
423 and argonaute 2 protein. *Non-Coding RNA* 2015, 1: 151-165

424 11. Trionfini P, Benigni A, Remuzzi G: MicroRNAs in kidney physiology and
425 disease. *Nat Rev Nephrol* 2015;11:23-33

426 12. Simpson K, Wonnacott A, Fraser DJ, Bowen T: MicroRNAs in diabetic
427 nephropathy: from biomarkers to therapy. *Curr Diab Rep* 2016, 16:35

428 13. Krupa A, Jenkins R, Luo DD, Lewis A, Phillips AO, Fraser DJ: Loss of MicroRNA-
429 192 promotes fibrogenesis in diabetic nephropathy. *J Am Soc Nephrol* 2010,
430 21:438-447

431 14. KDOQI Clinical Practice Guideline for Diabetes and CKD: 2012 Update. *Am J*
432 *Kidney Dis* 2012, 60:850-886

433 15. Stringer S, Sharma P, Dutton M, Jesky M, Ng K, Kaur O, Chapple I, Dietrich T,
434 Ferro C, Cockwell P: The natural history of, and risk factors for, progressive
435 chronic kidney disease (CKD): the Renal Impairment in Secondary care (RIISC)
436 study; rationale and protocol. *BMC Nephrol* 2013, 14:95

- 437 16. American Diabetes Association: Diagnosis and classification of diabetes
438 mellitus. Diabetes Care 2010, 33(suppl 1):S62-S69
- 439 17. Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D: A more accurate
440 method to estimate glomerular filtration rate from serum creatinine: a new
441 prediction equation. Modification of Diet in Renal Disease Study Group. Ann
442 Intern Med 1999, 130:461-470
- 443 18. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group (2013)
444 KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management
445 of Chronic Kidney Disease. Kidney Int Suppl 2013, 3:S1-150
- 446 19. Satchell SC, Tasman CH, Singh A, Ni L, Geelen J, von Rhuland CJ, O'Hare MJ,
447 Saleem MA, van den Heuvel LP, Mathieson PW: Conditionally immortalized
448 human glomerular endothelial cells expressing fenestration in response to
449 VEGF. Kidney Int 2006, 69:1633-1640
- 450 20. Saleem MA, O'Hare MJ, Reiser J, Coward RJ, Inward CD, Farren T, Xing CY, Ni L,
451 Mathieson PW, Mundel P: A conditionally immortalized human podocyte cell
452 line demonstrating nephrin and podocin expression. J Am Soc Nephrol 2002,
453 13:630-638
- 454 21. Jenkins RH, Davies LC, Taylor PR, Akiyama H, Cumbers B, Beltrami C,
455 Carrington CP, Phillips AO, Bowen T, Fraser DJ: MiR-192 induces G₂/M growth
456 arrest in aristolochic acid nephropathy. Am J Pathol 2014, 184:996-1009
- 457 22. Midgley AC, Bowen T, Phillips AO, Steadman R: MicroRNA-7 inhibition
458 rescues loss of epidermal growth factor receptor hyaluronan-dependent
459 differentiation in fibroblasts. Aging Cell 2014, 13:235-244

- 460 23. Zhang SD, Gant TW: Effect of pooling samples on the efficiency of
461 comparative studies using microarrays. *Bioinformatics* 2005, 21:4378-4383
- 462 24. Kendzierski C, Irizarry RA, Chen KS, Haag JD, Gould MN: On the utility of
463 pooling biological samples in microarray experiments. *Proc Natl Acad Sci USA*
464 2005, 102:4252-4257
- 465 25. Prowle JR, Calzavacca P, Licari E, Ligabo EV, Echeverri JE, Bagshaw SM,
466 Haase-Fielitz A, Haase M, Ostland V, Noiri E, Westerman M, Devarajan P,
467 Bellomo R: Combination of biomarkers for diagnosis of acute kidney injury
468 after cardiopulmonary bypass. *Ren Fail* 2015, 37:408–416, 2015
- 469 26. Schley G, Köberle C, Manuilova E, Rutz S, Forster F, Weyand M, Formentini I,
470 Kientsch-Engel R, Kai-Eckardt U, Willam C: Comparison of plasma and urine
471 biomarker performance in acute kidney injury. *PLoS ONE* 2015, 10:e0145042
- 472 27. Navarro J, Milena F, Mora C, Leon C, Claverie F, Flores C, Garcia J: Tumor
473 necrosis factor-alpha gene expression in diabetic nephropathy: Relationship
474 with urinary albumin excretion and effect of angiotensin-converting enzyme
475 inhibition. *Kidney Int Suppl* 2005, 99:S98 –S102
- 476 28. Ohga S, Shikata K, Yozai K, Okada S, Ogawa D, Usui H, Wada J, Shikata Y,
477 Makino H: Thiazolidinedione ameliorates renal injury in experimental diabetic
478 rats through anti-inflammatory effects mediated by inhibition of NF- κ B
479 activation. *Am J Physiol-Renal Physiol* 2007, 292:F1141–F1150
- 480 29. Smith DA, Newbury LJ, Drago G, Bowen T, Redman JE: Electrochemical
481 detection of urinary microRNAs via sulfonamide-bound antisense
482 hybridisation. *Sens Actuators B Chem* 2017, 253:335–341

- 483 30. Olivieri F, Spazzafumo L, Bonafè M, Recchioni R, Prattichizzo F, Marcheselli M,
484 Micolucci L, Mensà E, Giuliani A, Santini G, Gobbi M, Lazzarini R, Boemi M,
485 Testa R, Antonicelli R, Antonio Domenico Procopio AD, Bonfigli AR: MiR-21-5p
486 and miR-126a-3p levels in plasma and circulating angiogenic cells:
487 relationship with type 2 diabetes complications. *Oncotarget* 2015, 6:35372-
488 35382
- 489 31. Barutta F, Bruno G, Matullo G, Chaturvedi N, Grimaldi S, Schalkwijk C,
490 Stehouwer CD, Fuller JH, Gruden G: MicroRNA-126 and micro-/macrovascular
491 complications of type 1 diabetes in the EURODIAB prospective complications
492 study. *Acta Diabetol* 2017, 54:133–139
- 493 32. Al-Kafaji G, Al-Mahroos G, Al-Muhtaresh HA, Skrypnyk C, Sabry MA, Ramadan
494 AR: Decreased expression of circulating microRNA-126 in patients with type 2
495 diabetic nephropathy: a potential blood-based biomarker. *Exp Therapeut*
496 *Med* 2016, 12:815-822
- 497 33. Osipova J, Fischer D-C, Dangwal S, Volkmann I, Widera C, Schwarz K,
498 Lorenzen JM, Schreiver C, Jacoby U, Heimhalt M, Thum T, Haffner D:
499 Diabetes-Associated MicroRNAs in Pediatric Patients With Type 1 Diabetes
500 Mellitus: A Cross-Sectional Cohort Study. *J Clin Endocrinol Metab* 2014,
501 99:E1661–E1665
- 502 34. Wang S, Aurora AB, Johnson BA, Qi X, McAnally J, Hill JA, Richardson JA,
503 Bassel-Duby R, Olson EN: The endothelial-specific microRNA miR-126 governs
504 vascular integrity and angiogenesis. *Dev Cell* 2008, 15:261–271
- 505 35. Brosius FC, Coward RJ: Podocytes, signaling pathways, and vascular factors in
506 diabetic kidney disease. *Adv Chronic Kidney Dis* 2014, 21:304–310

- 507 36. Harris TA, Yamakuchi M, Ferlito M, Mendell JT, Lowenstein CJ: MicroRNA-126
508 regulates endothelial expression of vascular cell adhesion molecule 1. *Proc*
509 *Natl Acad Sci USA* 2008, 105:1516–1521
- 510 37. Ásgeirsdóttir SA, van Solingen C, Kurniati NF, Zwiers PJ, Heeringa P, van
511 Meurs M, Satchell SC, Saleem MA, Mathieson PW, Banas B, Kamps JA,
512 Rabelink TJ, van Zonneveld AJ, Molema G: MicroRNA-126 contributes to renal
513 microvascular heterogeneity of VCAM-1 protein expression in acute
514 inflammation. *Am J Physiol Renal Physiol* 2012, 302:F1630-1639
- 515 38. Huang Y, Liu Y, Li L, Su B, Yang L, Fan W, Yin Q, Chen L, Cui T, Zhang J, Lu Y,
516 Cheng J, Fu P, Liu F: Involvement of inflammation-related miR-155 and miR-
517 146a in diabetic nephropathy: implications for glomerular endothelial injury.
518 *BMC Nephrol* 2014, 15:142
- 519 39. Lin X, You Y, Wang J, Qin Y, Huang P, Yang F: MicroRNA-155 deficiency
520 promotes nephrin acetylation and attenuates renal damage in
521 hyperglycemia-induced nephropathy. *Inflammation* 2015, 38:546-544
- 522 40. Welten SMJ, Goossens EAC, Quax PHA, Nossent AY: The multifactorial nature
523 of microRNAs in vascular remodeling. *Cardiovasc Res* 2016, 110:6–22
- 524 41. Wang B, Komers R, Carew R, Winbanks CE, Xu B, Herman-Edelstein M, Koh P,
525 Thomas M, Jandeleit-Dahm K, Gregorevic P, Cooper ME, Kantharidis P:
526 Suppression of microRNA-29 expression by TGF- β 1 promotes collagen
527 expression and renal fibrosis. *J Am Soc Nephrol* 23: 252-265, 2012
- 528 42. Chen H-Y, Zhong X, Huang XR, Meng X-M, You Y, Chung ACK, Lan HY:
529 MicroRNA-29b inhibits diabetic nephropathy in db/db Mice. *Mol Ther* 2014,
530 22:842-853

- 531 43. Toyono T, Usui T, Villarreal G, Kallay L, Matthaei M, Vianna LMM, Zhu AY,
532 Kuroda M, Amano S, Jun AS: MicroRNA-29b overexpression decreases
533 extracellular matrix mRNA and protein production in human corneal
534 endothelial cells. *Cornea* 2016, 35:1466–1470
- 535 44. Zhu HQ, Li Q, Dong LY, Zhou Q, Wang H, Wang Y: MicroRNA-29b promotes
536 high-fat diet-stimulated endothelial permeability and apoptosis in apoE
537 knock-out mice by down-regulating MT1 expression. *Int J Cardiol* 2014,
538 176:764-770
- 539 45. Silambarasan M, Tan JR, Karolina DS, Armugam A, Kaur C, Jeyaseelan K:
540 MicroRNAs in hyperglycemia induced endothelial cell dysfunction. *Int J Mol*
541 *Sci* 2016, 17:518
- 542 46. Mocharla P, Briand S, Giannotti G, Dörries C, Jakob P, Paneni F, Lüscher T,
543 Landmesser U: AngiomiR-126 expression and secretion from circulating
544 CD34⁺ and CD14⁺ PBMCs: role for proangiogenic effects and alterations in
545 type 2 diabetics. *Blood* 2013, 121:226-236
- 546 47. Barutta F, Tricarico M, Corbelli A, Annaratone L, Pinach S, Grimaldi S, Bruno
547 G, Cimino D, Taverna D, Deregibus MC, Rastaldi MP, Perin PC, Gruden G:
548 Urinary exosomal microRNAs in incipient diabetic nephropathy. *PLoS ONE*
549 2013, 8:e73798
- 550 48. Salama A, Fichou N, Allard M, Dubreil L, De Beaurepaire L, Viel A, Jégou D,
551 Bösch S, Bach J-M: MicroRNA-29b modulates innate and antigen-specific
552 immune responses in mouse models of autoimmunity. *PLoS ONE* 2014,
553 9:e106153

Figure legends

Figure 1 Urinary miRNA detection in urine samples from DKD patients and control subjects. **A:** Volcano plot showing the detection profile of the 377 urinary miRNAs in TLDA Card A in DKD patients (n = 20; four pools of five patients) and controls (n = 20; four pools of five controls). The dotted horizontal line represents a p value boundary of 0.05. **B:** Fold change of miRNA detection between DKD patients and controls. DataAssist Software (Thermo Fisher Scientific) was used to perform relative quantification for sample comparison, to perform t-test sample group comparisons, and to produce the graphic output shown. **(C-E)** RT-qPCR analysis shows significant differences in detection of **(C)** miR-126, **(D)** miR-155 and **(E)** miR-29b between patients and control urine in the component urine samples pooled for profiling analyses **(A and B)**. DKD patients versus controls for **C:** miR-126, **D:** miR-155, **E:** miR-29b (n = 20 for each group). Analysis was carried out by unpaired 2-tailed t-test with Welch's correction. Profiling data analysis using the NormFinder algorithm identified miR-191 as optimal for normalisation of our RT-qPCR data. Data were normalized to endogenous control miR-191 and are presented as mean +/- SEM. ** $P < 0.01$, *** $P < 0.005$ and **** $P < 0.0005$.

Figure 2 RT-qPCR detection of selected miRNAs in patients and control subjects. Relative expression was significantly different in 89 DKD patients compared with 62 diabetic patients without DKD and 41 controls for **(A)** miR-126 and **(B)** miR-155, and significantly different in DKD patients compared with controls for **(C)** miR-29b. **A:** DKD patients versus diabetic patients without DKD and controls; **B:** DKD patients and diabetic patients without DKD; DKD patients versus controls; **C:** DKD patients versus

controls (n = 89 DKD patients, 62 diabetic patients without DKD, 41 controls). Analysis was carried out by unpaired 1-tailed t-test with Welch's correction. Data were normalized to endogenous control miR-191 and are presented as mean +/- SEM. (D) Combined receiver operating characteristic curve analysis for miR-126, miR-155 and miR-29b, area under the curve (AUC) = 0.80. Data were generated using the pROC package in R-3.2.3. * $P < 0.05$, **** $P < 0.001$ and ***** $P < 0.0001$.

Figure 3 Localization of miRNA expression by laser capture microdissection (LCM) and cell culture. **A:** Key functional nephron domains include the glomerulus (G), the proximal tubule (PT) and the distal tubule (DT). **B:** A CD10-stained FFPE renal biopsy sample before and after excision of glomeruli by LCM. Bars = 100 μ m. **(C-E)** Relative expression of miR-126, miR-155 and miR-29b, respectively, in LCM-isolated Gs, PTs and DTs from 5 renal biopsies of healthy individuals. **C:** G versus PT; G versus DT (n = 5 biopsies). **(F-H)** Relative expression of miR-126, miR-155 and miR-29b, respectively, in *in vitro* cultured HK-2 renal proximal tubular epithelial cells (PTCs), fibroblasts, podocytes and conditionally immortalized glomerular endothelial cells (GEnCs). **F:** GEnCs versus fibroblasts; GEnCs versus PTCs and podocytes; **G:** PTC versus GEnCs; **H:** GEnCs versus podocytes (n = 4). Analysis was carried out by one-way ANOVA analysis with Tukey's multiple comparison test. Data were normalized to endogenous control miR-191 and are presented as mean +/- SEM. * $P < 0.05$, ** $P < 0.01$ and **** $P < 0.001$.

Figure 4 MiRNA expression in GenCs and GenC conditioned medium in response to hyperglycaemia and DKD-related cytokines. Following 24 h culture in 5 mM or 25 mM D-glucose, relative expression in GEnCs and GEnC conditioned medium, respectively, of **(A and B)** miR-126 in response to 10 ng/ml TNF- α , **(C and D)** miR-29b in response to 1 ng/ml TGF- β 1 and **(E and F)** miR-155 in response to 10 ng/ml TNF- α , and in untreated cells. **B:** 5 mM D-glucose versus 5 mM D-glucose plus TNF- α , and 25 mM D-glucose versus 25 mM D-glucose plus TNF- α ; **C:** 5 mM D-glucose versus 5 mM D-glucose plus TGF- β 1, and 25 mM D-glucose versus 25 mM D-glucose plus TGF- β 1; **D:** 5 mM D-glucose versus 5 mM D-glucose plus TGF- β 1; 25 mM D-glucose versus 25 mM D-glucose plus TGF- β 1 (n = 4). Analysis was carried out by one-way ANOVA analysis with Tukey's multiple comparison test. Data were normalized to endogenous control miR-191 and are presented as mean \pm SEM. * P < 0.05 and ** P < 0.01.

Table 1 Demographic and clinical parameters of patients recruited from 2 centres:
Wales Kidney Research Tissue Bank, Cardiff (University Hospital Wales) and
Birmingham (University Hospital Birmingham, Renal Impairment in Secondary care
(RIISC) study cohort) (n = 151)

Feature	Patients (n=151)		Controls (n=41)
	Diabetic (n=62)	DKD (n=89)	
Male n (%)	37 (58)	55 (62)	18 (44)
Non-Caucasian n (%)	13 (21)	33 (37)	
Mean Age (years, SD)	52 +/- 16.1	62 +/- 13.6	55 +/- 15.4
eGFR ml/min/1.73m ²			
Mean (SD)	78 +/-16.3	30 +/- 20.9	
Median (IQR)	84 (72-90)	22 (17-38)	
CKD stage n(%)			
No CKD/CKD G1 (eGFR ≥ 90)	23 (37)	2 (2)	
CKD G2 (eGFR = 60-89)	32 (52)	10 (11)	
CKD G3 (eGFR = 30-59)	5 (8)	17 (19)	
CKD G4 (eGFR = 15-29)	2 (3)	45 (51)	
CKD G5 (eGFR <15)	0	15 (17)	
ACR* mg/mmol n(%)			
A1-Normal-high normal (ACR<3)	54 (87.1)	15 (16.9)	
A2-Moderately increased (ACR 3-30)	8 (12.9)	25 (28.1)	
A3-Severely increased (ACR>30)	0 (0)	49 (55.0)	

*Albumin:Creatinine ratio (ACR) group cut-offs and nomenclature derived from
KDIGO 2012 recommendations.

Table 2 Specificity values, likelihood ratios and RQ thresholds for miR-126, miR-155, miR-29b and all three miRNAs above an 80% ROC curve sensitivity threshold

	Sensitivity (%)	Specificity (%)	Likelihood Ratio	RQ Threshold
3 miRNAs	80.21	63.64	2.206	> 1.148
miR-126	80.41	57.14	1.876	> 0.6762
miR-155	80.61	52.00	1.679	> 0.9110
miR-29b	80.61	40.00	1.344	> 0.8058

Table 3 DKD and diabetic patients without DKD (D) patient numbers and percentages above an 80% ROC curve sensitivity threshold for miR-126, miR-155, miR-29b and all three miRNAs

	Patients above 80% sensitivity threshold				Total
	miR-126	miR-155	miR-29b	3 miRNAs	
D	23	30	13	2	55
DKD	80	67	73	47	98

	Percentage of patients above 80% sensitivity threshold			
	miR-126	miR-155	miR-29b	3 miRNAs
D	41.8	54.6	23.6	3.6
DKD	81.6	68.4	74.5	48.0




Figure 1







Figure 2



B






Renal biopsy before LCM



Renal biopsy following LCM

LCM-dissected
nephron domains



Cell cultures






Figure 3




Figure 4